

A new attraction-detachment model for explaining flow sliding in clay-rich tephtras

Max O. Kluger¹, Vicki G. Moon², Stefan Kreiter¹, David J. Lowe², G.J. Churchman³, Daniel A. Hepp¹, David Seibel¹, M. Ehsan Jorat⁴, and Tobias Mörz¹

¹MARUM—Center for Marine Environmental Sciences, University of Bremen, Leobener Straße, 28359 Bremen, Germany

²School of Science, University of Waikato, Private Bag 3105, Hamilton 3240, New Zealand

³School of Agriculture, Food and Wine, University of Adelaide, Adelaide, SA 5005, Australia

⁴School of Science, Engineering and Technology, Abertay University, Dundee DD11HG, UK

ABSTRACT

Altered pyroclastic (tephra) deposits are highly susceptible to landsliding, leading to fatalities and property damage every year. Halloysite, a low-activity clay mineral, is commonly associated with landslide-prone layers within altered tephra successions, especially in deposits with high sensitivity, which describes the post-failure strength loss. However, the precise role of halloysite in the development of sensitivity, and thus in sudden and unpredictable landsliding, is unknown. Here we show that an abundance of mushroom cap-shaped (MCS) spheroidal halloysite governs the development of sensitivity, and hence proneness to landsliding, in altered rhyolitic tephtras, North Island, New Zealand. We found that a highly sensitive layer, which was involved in a flow slide, has a remarkably high content of aggregated MCS spheroids with substantial openings on one side. We suggest that short-range electrostatic and van der Waals interactions enabled the MCS spheroids to form interconnected aggregates by attraction between the edges of numerous paired silanol and aluminol sheets that are exposed in the openings and the convex silanol faces on the exterior surfaces of adjacent MCS spheroids. If these weak attractions are overcome during slope failure, multiple, weakly attracted MCS spheroids can be separated from one another, and the prevailing repulsion between exterior MCS surfaces results in a low remolded shear strength, a high sensitivity, and a high propensity for flow sliding. The evidence indicates that the attraction-detachment model explains the high sensitivity and contributes to an improved understanding of the mechanisms of flow sliding in sensitive, altered tephtras rich in spheroidal halloysite.

INTRODUCTION

Most East Asian and western Pacific countries are located in tectonically active, high-rainfall areas where landslides are a major natural hazard. These landslides are typically triggered by rainstorms or earthquakes and are responsible for fatalities and enormous property damage every year. Many destructive landslides have occurred in pyroclastic deposits in Japan, Indonesia, Hong Kong, and New Zealand (Chau et al., 2004; Chigira, 2014; Moon, 2016), such deposits commonly containing layers rich in clay minerals formed mainly by chemical weathering during either pedogenesis or diagenesis. In regions with predominantly rhyolitic volcanism, halloysite is a common clay mineral (Churchman and Lowe, 2012) and is therefore potentially a key geological factor increasing the risk of landslides (Kirk et al., 1997; Chigira, 2014). Halloysite is a 1:1 Si:Al layered aluminosilicate member of the kaolin subgroup that exhibits various structural morphologies including tubes, spheroids, polyhedrons, plates, and books (Joussein et al., 2005; Cunningham et al., 2016).

Spheroidal halloysite, in particular, has been recognized in landslide-prone layers of pyroclastic material in Japan (Tanaka, 1992) and New Zealand (Smalley et al., 1980). Smalley et al. (1980) linked a high content of spheroidal halloysite to high sensitivity. Sensitivity refers to the post-failure strength loss in the failure zone during landsliding, and is quantified in the laboratory as the ratio of the undisturbed to remolded undrained shear strength at the same water content (Terzaghi, 1944). High sensitivities were first described for post-glacial, brackish and marine clayey sediments in the Northern Hemisphere (Skempton and Northey, 1952) that are subject to landslides with dimensions and long runout distances that are difficult to predict. In this study, we investigate processes that have led to high sensitivity in halloysite-rich pyroclastic materials in order to improve landslide-hazard evaluation.

GEOLOGICAL SETTING

Much of the central part of New Zealand's North Island is covered by thick rhyolitic tephtras (Lowe, 2011) derived from eruptions in

the Taupo Volcanic Zone (Briggs et al., 2005), which are commonly altered into halloysite-rich successions. We focus here on a coastal flow slide at Omokoroa, Bay of Plenty (Fig. 1A), where ~10,000 m³ of material was transported downslope over long distance into a lagoon in 1979 CE (Moon et al., 2015) as well as during two minor reactivations in 2011 and 2012. The 1979 event was likely initiated in a white, highly sensitive layer with high spheroidal halloysite concentration (Smalley et al., 1980) (and lacking any detectable allophane; Cunningham et al., 2016).

We have analyzed a 40-m-long sediment core, Omok-1, which we bored via rotary flush drilling in unfailed material near the headwall (Fig. 1B). The lithology of Omok-1 was determined by correlation with units of a previously studied adjacent headwall face (Moon et al., 2015) comprising a succession mainly of Quaternary rhyolitic tephtras: underlain by lignite at the base of the core, the Pahoia Tephra sequence includes the Te Puna Ignimbrite (ca. 0.93 Ma) and a series of altered tephtras which are informally divided into lower and upper Pahoia Tephra units based on two distinct paleosols (P1 and P3). All of these deposits and paleosols are overlain by successions of younger altered tephtras called Hamilton Ash beds (ca. 0.35 to ca. 0.05 Ma) and late Quaternary tephtras (< ca. 0.05 Ma) (Figs. 1C and 2A). The lower Pahoia Tephtras include the 0.3-m-thick, white, highly sensitive clay-rich layer that failed in 1979 (Fig. 1C), having high porosity and high natural water content (Smalley et al., 1980).

METHODS

We performed laboratory vane shear tests on samples from the Pahoia Tephra sequence and Hamilton Ash beds to measure the sensitivity S :

$$S = s_u / s_r, \quad (1)$$

where the undisturbed strength (s_u) was measured on the intact surface of the split core, and the remolded strength (s_r) was measured on core




Figure 1. A: Map of Tauranga Harbour, New Zealand, with Taupo Volcanic Zone (TVZ) as main source for Quaternary tephras at study site. **B:** Three-dimensional view of the flow slide at Omokoroa; red line marks position of profile in C. **I.**—Ignimbrite. **C:** Profile through flow slide with simplified stratigraphy and associated paleosols (P1–P4) of drill core Omok-1 and ages (in Ma) after Moon et al. (2015).

samples with the same water content but that had been kneaded by hand for 10 min (Jacquet, 1990). Halloysite concentration in bulk samples was measured by X-ray diffraction (XRD) using a Philips PW analytical diffractometer,

and quantification was performed using QUAX software (Vogt et al., 2002). Scanning electron microscopy (SEM) was undertaken with a Zeiss Supra40 microscope on 24 shock-frozen, freeze-dried, and gold-coated bulk core samples

(Reed, 2005). The relative abundances of halloysite particles having distinct morphologies were quantified using a point-counting approach (Frolov and Maling, 1969). Six representative SEM images of planar soil surfaces were chosen for each sample, and at least 600 particles were counted based on rectangular grids. In the white, highly sensitive layer, the change of halloysite particle arrangement upon remolding was quantified by comparing 20 SEM images of undisturbed and remolded material, providing >1000 counts respectively. The spheroid diameters were measured from six representative particles per SEM image.

HIGHLY SENSITIVE SLIDE-PRONE LAYER DOMINATED BY SPHEROIDAL HALLOYSITE

The sensitivity is low in the upper Pahoia Tephra, especially in the paleosols P2 and P3 (Figs. 2A and 2B). However, the sensitivity tends to increase with depth, reaching values of 15–20 in the lower Pahoia Tephra. The highest sensitivity (Rosenqvist, 1953) of $S = 55$, and the lowest remolded shear strength within the profile of $s_r = 1.4$ kPa, were measured in the white, highly sensitive layer at 23 m depth.

The upper Pahoia Tephra have a halloysite content of 10–20 wt% comprising almost entirely tubular halloysite (Figs. 2C and 2D). The lower Pahoia Tephra have 40–50 wt% halloysite comprising mostly spheroidal particles. In the highly sensitive layer, 76% of the halloysite is spheroidal and the spheroid sizes




Figure 2. A: Stratigraphy of drill core Omok-1 after Moon et al. (2015) showing main lithological units as defined in Figure 1, three paleosols (P1–P3), and highly sensitive white layer at 23 m depth (hatched area). **I.**—Ignimbrite; **T.**—Tephra. **B:** Undisturbed (s_u , blue) and remolded (s_r , orange) shear strength, and sensitivity ($S = s_r / s_u$). **C:** Halloysite bulk concentration. **D:** Cumulative volume percent (c. vol%) of halloysite morphologies with bars indicating average standard deviations. **E:** Average spheroid sizes with standard deviations depicted by fill patterns. **F:** Three-dimensional line plot illustrating relationship between spheroid content, sensitivity, spheroid size, and halloysite concentration; gray graded areas enable trends in sensitivity to be visualized. **G:** Dependency between sensitivity and shear strength.

are greater than those in the surrounding layers (Figs. 2D and 2E). A three-dimensional line plot reveals a clear correlation between high sensitivities and high halloysite bulk concentration, and a high content of spheroids with large diameters (Fig. 2F). The high sensitivity is associated with low remolded shear strength rather than with high undisturbed shear strength (Fig. 2G).

We found that deposits with high tubular halloysite content hamper sensitivity development, whereas halloysite spheroids facilitate sensitivity and dominate the highly sensitive layer at 23 m depth within the lower Pahoia Tephra. The highly sensitive layer has low remolded shear strength after failure, which, together with its high water content (Smalley et al., 1980), partly contributed to the long runout distance of the flow slide at Omokoroa.

NEW HALLOYSITE MORPHOLOGY

We present here the first observations of a previously unreported halloysite particle morphology, which is visible in the SEM images of the remolded halloysite fabrics of the highly sensitive layer. In the undisturbed state, the spheroidal halloysites are distinctly aggregated into networks of well-connected particles (Figs. 3E and 3F). After remolding, however, most of the aggregates have broken apart into small, loose clusters or individual halloysite particles that are typically ~250–400 nm in diameter (Figs. 3G and 3H). Individual spheroids have distinctive “deformities” in the form of openings ~80–160 nm in diameter on one side. These openings were previously hidden by contact with other spheroids. The deformities give the particles an ovate “mushroom cap” appearance. Point-counting individual mushroom-cap shapes in both undisturbed (aggregated) and remolded (disaggregated) samples showed that the observable mushroom-cap shapes were much more abundant in the remolded samples, increasing from $4.4\% \pm 3.2\%$ to $44.9\% \pm 11.6\%$.

ATTRACTION-DETACHMENT MODEL FOR FLOW SLIDING IN ALTERED TEPHRAS

The open-sided, mushroom cap-shaped halloysite morphology has not been reported previously. Because this particular morphology overwhelmingly occurs in the highly sensitive slide-prone layer, we hypothesize that this unique particle shape controls the mechanical behavior of halloysite clays.

Halloysite is composed of an Al-octahedral (aluminol) sheet with a net positive charge and a Si-tetrahedral (silanol) sheet with a net negative charge at pH values between ~2 and ~8 (Fig. 3I) (Churchman et al., 2016). The two sheets have slightly different dimensions, with the silanol sheet being larger. This misfit in the sheet sizes causes the halloysite layer to be curved (Churchman and Lowe, 2012), with the larger negatively




Figure 3. A–D: Scanning electron microscopy (SEM) images of spheroids (A), polyhedrons (B), tubes (C), and plates (D) representing the main halloysite morphologies in Pahoia Tephra sequence (New Zealand). E–H: SEM images from the highly sensitive layer of undisturbed and multiply-connected halloysite spheroids (E, F) and remolded spheroids (G, H) showing smaller clusters or detached spheroids within much looser particle network. 1—exposed layers in spheroid openings; 2—partially separated halloysite spheroids; 3—detached mushroom cap-shaped halloysite spheroid. I: Electrostatic field proximal to halloysite nanotubes with colored equipotential surfaces (ES), modified with permission from Guimarães et al. (2010), copyright 2010 American Chemical Society. J: Conceptual mushroom cap-shaped spheroid cross-section and weak electrostatic and/or van der Waals attractions arising between exposed silanol-aluminol sheets in spheroid openings and negatively charged convex exterior surfaces; enlargement is adapted from Berthonneau et al. (2015). Circles with + and – relate to positive and negative electrostatic field proximal to spheroid’s exterior surface. K, L: Mushroom cap-shaped spheroids connect with one another between concave openings and convex outer spheroid surfaces, forming aggregates (K) which are partly detached because of remolding (L).

charged silanol sheet on the outside of the curvature and the positively charged smaller aluminol sheet on the inside. The halloysite spheroids observed in our study are most likely composed of concentrically stacked 1:1 layers, i.e., with an onion-like structure, as shown in numerous studies including those on spheroidal halloysite derived from altered tephra in New Zealand, Japan, and Argentina (Wada et al., 1977; Kirkman, 1981; Cravero et al., 2012; Berthonneau

et al., 2015). For a perfect halloysite spheroid, the outermost silanol surface carries a net negative charge, and hence the electrostatic interactions between individual spheroids would be repulsive (Fig. 3I). Our study shows, however, a halloysite structure where both silanol and aluminol layers are exposed at spheroid openings, and therefore charges within the openings would correspondingly be weakly positive or neutral overall (Fig. 3J), as indicated from

charge density-functional tight-binding modeling applied to halloysite nanotubes (Guimarães et al., 2010). If sufficient numbers of positively charged openings are exposed, the electrostatic interactions between them and the negative exterior silanol surfaces would allow the mushroom cap-shaped spheroids to form stacked aggregates (Fig. 3K). If the paired silanol and aluminol sheets exposed in the openings are neutral overall, then a net increase in particle attraction will still occur because electrostatic repulsion is reduced and the larger contact areas lead to higher van der Waals forces (Israelachvili, 2011).

During diagenesis via hydrolysis of volcanic glass (Cunningham et al., 2016), the halloysite spheroids may form consecutively on top of one another in pore spaces, generating the distinct openings during synthesis. The attractive forces between the openings and the convex exterior surfaces are demonstrably strong enough to allow for the formation of aggregates, but also permit easy disaggregation by mechanical detachment during shear (Fig. 3L). New random contacts between convex silanol surfaces probably lead to a decrease in average attraction between particles. We posit that the detachment of attractive spheroidal particle contacts, in the presence of abundant water having negligible interaction with ions in soil solution because of the inactive nature of halloysite (Smalley et al., 1980), leads to the very low post-failure shear strength, facilitating a flow slide with long runout distance. The interparticle attraction-detachment model appears to successfully explain (at nanoscale dimensions) the post-failure behavior of the highly sensitive tephra layer at Omokoroa, which is dominated by the imperfect halloysite spheroids. The question therefore arises whether similar altered tephtras elsewhere have high contents of spheroidal halloysite with potentially hidden mushroom-cap forms, and whether such forms helped mobilize other landslides in the past.

CONCLUSIONS

We investigated a sequence of altered, rhyolitic Quaternary tephtras in New Zealand and the reasons why a landslide-prone layer dominated by spheroidal halloysite was highly sensitive. We explain this high sensitivity with an electrostatic attraction-detachment model. Weakly positive or neutral charges on silanol and aluminol sheet edges exposed in the concave openings of spheroidal halloysite particles were attracted to the negatively charged convex silanol surfaces of adjacent spheroids. Such short-range attractions between spheroid openings and the exterior surfaces of adjacent spheroids stabilize an aggregated halloysite framework. If the aggregates are detached by remolding, the loose arrangement of the spheroids exhibits low remolded

shear strength. We suggest that the attraction-detachment model, based on the identification of mushroom-cap halloysite morphologies, provides a potential key for the identification of sensitive altered tephtras that are predisposed to sudden failure that triggers landsliding.

ACKNOWLEDGMENTS

This research was funded by the DFG Research Center MARUM (University of Bremen) through INTERCOAST and the University of Waikato (Hamilton, New Zealand). We thank C. Schulze for vane shear tests; B. Steinborn and C. Vogt for XRD analyses; P. Witte, A. Hübner, C. Schott, S. Buchheister, and V. Diekamp for laboratory assistance; M. Ikari, F. Sense, J. Lane, and P. Pasbakhsh for comments; and F. Terrible, J.K. Torrance, F. Cravero, M. McSaveney, and an anonymous reviewer for helpful reviews.

REFERENCES CITED

- Berthonneau, J., Grauby, O., Jeannin, C., Chaudanson, D., Joussein, E., and Baronnet, A., 2015, Native morphology of hydrated spheroidal halloysite observed by environmental transmission electron microscopy: *Clays and Clay Minerals*, v. 63, p. 368–377, doi:10.1346/CCMN.2015.0630503.
- Briggs, R., Houghton, B., McWilliams, M., and Wilson, C., 2005, ⁴⁰Ar/³⁹Ar ages of silicic volcanic rocks in the Tauranga-Kaimai area, New Zealand: Dating the transition between volcanism in the Coromandel Arc and the Taupo Volcanic Zone: *New Zealand Journal of Geology and Geophysics*, v. 48, p. 459–469, doi:10.1080/00288306.2005.9515126.
- Chau, K., Sze, Y., Fung, M., Wong, W., Fong, E., and Chan, L., 2004, Landslide hazard analysis for Hong Kong using landslide inventory and GIS: *Computers & Geosciences*, v. 30, p. 429–443, doi:10.1016/j.cageo.2003.08.013.
- Chigira, M., 2014, Geological and geomorphological features of deep-seated catastrophic landslides in tectonically active regions of Asia and implications for hazard mapping: *Episodes*, v. 37, p. 284–294.
- Churchman, G.J., and Lowe, D.J., 2012, Alteration, formation, and occurrence of minerals in soils, in Huang, P.M., et al., eds., *Handbook of Soil Sciences: Properties and Processes* (second edition): Boca Raton, Florida, CRC Press, p. 20.21–20.72.
- Churchman, G.J., Pasbakhsh, P., Lowe, D.J., and Theng, B.K.G., 2016, Unique but diverse: Some observations on the formation, structure, and morphology of halloysite: *Clay Minerals*, v. 51, p. 395–416, doi:10.1180/claymin.2016.051.3.14.
- Cravero, F., Maiza, P., and Marfil, S., 2012, Halloysite in Argentinian deposits: Origin and textural constraints: *Clay Minerals*, v. 47, p. 329–340, doi:10.1180/claymin.2012.047.3.04.
- Cunningham, M.J., Lowe, D.J., Wyatt, J.B., Moon, V.G., and Churchman, G.J., 2016, Discovery of halloysite books in altered silicic Quaternary tephtras, northern New Zealand: *Clay Minerals*, v. 51, p. 351–372, doi:10.1180/claymin.2016.051.3.16.
- Frolov, Y., and Maling, D., 1969, The accuracy of area measurement by point counting techniques: *The Cartographic Journal*, v. 6, p. 21–35, doi:10.1179/caj.1969.6.1.21.
- Guimarães, L., Enyashin, A.N., Seifert, G., and Duarte, H.A., 2010, Structural, electronic, and mechanical properties of single-walled halloysite nanotube models: *The Journal of Physical*

- Chemistry C*, v. 114, p. 11,358–11,363, doi:10.1021/jp100902e.
- Israelachvili, J.N., 2011, *Intermolecular and Surface Forces* (third edition): Waltham, Massachusetts, Academic Press, 704 p.
- Jacquet, D., 1990, Sensitivity to remoulding of some volcanic ash soils in New Zealand: *Engineering Geology*, v. 28, p. 1–25, doi:10.1016/0013-7952(90)90031-U.
- Joussein, E., Petit, S., Churchman, J., Theng, B., Righi, D., and Delvaux, B., 2005, Halloysite clay minerals—A review: *Clay Minerals*, v. 40, p. 383–426, doi:10.1180/00098550504040180.
- Kirk, P., Campbell, S., Fletcher, C., and Merriman, R., 1997, The significance of primary volcanic fabrics and clay distribution in landslides in Hong Kong: *Journal of the Geological Society*, v. 154, p. 1009–1019, doi:10.1144/gsjgs.154.6.1009.
- Kirkman, J.H., 1981, Morphology and structure of halloysite in New Zealand tephtras: *Clays and Clay Minerals*, v. 29, p. 1–9, doi:10.1346/CCMN.1981.0290101.
- Lowe, D.J., 2011, Tephrochronology and its application: A review: *Quaternary Geochronology*, v. 6, p. 107–153, doi:10.1016/j.quageo.2010.08.003.
- Moon, V., 2016, Halloysite behaving badly: Geomechanics and slope behaviour of halloysite-rich soils: *Clay Minerals*, v. 51, p. 517–528, doi:10.1180/claymin.2016.051.3.09.
- Moon, V.G., Lowe, D.J., Cunningham, M.J., Wyatt, J.B., de Lange, W.P., Churchman, G.J., Mörz, T., Kreiter, S., Kluger, M.O., and Jorat, M.E., 2015, Sensitive pyroclastic-derived halloysitic soils in northern New Zealand: Interplay of microstructure, minerals, and geomechanics, in Rotonda, T., et al., eds., *Volcanic Rocks and Soils: Proceedings of the International Workshop on Volcanic Rocks and Soils*, Lacco Ameno, Ischia Island, Italy: London, Taylor and Francis, p. 3–21.
- Reed, S.J.B., 2005, *Electron Microprobe Analysis and Scanning Electron Microscopy in Geology*: Cambridge, UK, Cambridge University Press, 212 p., doi:10.1017/CBO9780511610561.
- Rosenqvist, I.T., 1953, Considerations on the sensitivity of Norwegian quick-clays: *Géotechnique*, v. 3, p. 195–200, doi:10.1680/geot.1953.3.5.195.
- Skempton, A., and Northey, R., 1952, The sensitivity of clays: *Géotechnique*, v. 3, p. 30–53, doi:10.1680/geot.1952.3.1.30.
- Smalley, I., Ross, C.W., and Whitton, J., 1980, Clays from New Zealand support the inactive particle theory of soil sensitivity: *Nature*, v. 288, p. 576–577, doi:10.1038/288576a0.
- Tanaka, K., 1992, Slope hazards and clay minerals [in Japanese]: *Nendo Kagaku (Journal of the Clay Science Society of Japan)*, v. 32, p. 16–22, doi:10.11362/jcssjnedokagaku1961.32.16.
- Terzaghi, K., 1944, Ends and means in soil mechanics: *Engineering Journal of Canada*, v. 27, p. 608–615.
- Vogt, C., Lauterjung, J., and Fischer, R.X., 2002, Investigation of the clay fraction (<2 μm) of the Clay Minerals Society reference clays: *Clays and Clay Minerals*, v. 50, p. 388–400, doi:10.1346/000986002760833765.
- Wada, S.-I., Aoki, K., and Wada, K., 1977, The interior structure of spherical halloysite particles: *Clay Science*, v. 5, p. 113–121, doi:10.11362/jcssjclayscience1960.5.113.

Manuscript received 1 September 2016

Revised manuscript received 31 October 2016

Manuscript accepted 1 November 2016

Printed in USA